



Polynesian Journal of Mathematics

Volume 2, Issue 5

**On a minimum fractional
Hardy–Hilbert-type integral inequality**

Christophe Chesneau

Received 29 Jul 2025

Revised 3 Sep 2025

Accepted 5 Sep 2025

Published 17 Sep 2025

Communicated by Gaetan Bisson

DOI: 10.69763/polyjmath.2.5

On a minimum fractional Hardy–Hilbert-type integral inequality

Christophe Chesneau

Department of Mathematics, LMNO, University of Caen Normandie, France

Abstract

This article is devoted to the study of a new minimum fractional Hardy–Hilbert integral inequality. A comprehensive theoretical framework is developed, accompanied by a detailed proof. In addition, two auxiliary results are presented to illustrate the applicability and relevance of the main theorem.

Keywords: Integral inequalities, Hardy–Hilbert-type integral inequalities.

1 Introduction

Hardy–Hilbert-type integral inequalities have long occupied a central place in mathematical analysis. They play a particularly important role in the study of integral operators and functional inequalities. They also feature in numerous applications in the fields of harmonic analysis and partial differential equations. Since the pioneering work of Hardy, Hilbert and their contemporaries, these inequalities have been the subject of extensive generalizations and refinements, giving rise to a rich and continually evolving theoretical framework. Further information can be found in the books [6, 20, 21, 22], supplemented by the survey [4].

Considerable effort has recently been devoted to developing new variants of Hardy–Hilbert-type integral inequalities involving fractional, weighted and multilinear structures. These generalizations broaden the scope of classical inequalities. Among these developments, maximum- and minimum-type extensions have attracted particular attention due to their potential applications in the study of analytic inequalities, operator theory, and special functions. Notable contributions can be found in [19, 8, 9, 1, 13, 15, 17, 18, 16, 14, 5, 2, 11, 7, 3, 12, 10].

This article focuses on establishing a new minimum fractional Hardy–Hilbert-type integral inequality. Our approach builds upon and extends existing techniques, incorporating a more detailed analysis of parameter ranges and integrability conditions. The main contribution of this article is the formulation and proof of a sharp inequality involving a double integral and a minimum function of the ratios x/y and y/x , an

exponent parameter, α , and two main functions, f and g , i.e.,

$$\int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x)g(y)dx dy.$$

Based on this result, we further derive several secondary, yet new, integral inequalities.

The remainder of the article is organized as follows. Section 2 presents the main theorem. In Section 3, we develop the secondary results. Finally, Section 4 provides concluding remarks and outlines possible directions for future research.

2 Main result

Our main result is the theorem stated below.

Theorem 2.1. *Let $p > 1$, $q = p/(p - 1)$, $\alpha, \beta > 0$ such that $\alpha > \max(|\beta p - 1|, |\beta q - 1|)$, and $f, g : (0, +\infty) \mapsto (0, +\infty)$ be two functions. Then we have*

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x)g(y)dx dy \\ & \leq \frac{2\alpha}{(\alpha - \beta p + 1)^{1/p}(\alpha + \beta p - 1)^{1/p}(\alpha - \beta q + 1)^{1/q}(\alpha + \beta q - 1)^{1/q}} \\ & \quad \times \left[\int_{(0,+\infty)} x f^p(x) dx \right]^{1/p} \left[\int_{(0,+\infty)} y g^q(y) dy \right]^{1/q} \end{aligned}$$

provided that the two integrals involved in the upper bound converge.

Proof. By suitably decomposing the integrand via the identity

$$1 = \left(\frac{x}{y} \right)^\beta \left(\frac{y}{x} \right)^\beta$$

and applying the Hölder integral inequality with exponents satisfying $1/p + 1/q = 1$, we obtain

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x)g(y)dx dy \\ & = \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^{\alpha/p} \left(\frac{x}{y} \right)^\beta f(x) \\ & \quad \times \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^{\alpha/q} \left(\frac{y}{x} \right)^\beta g(y)dx dy \\ & \leq \mathfrak{A}^{1/p} \mathfrak{B}^{1/q} \end{aligned} \tag{1}$$

where \mathfrak{A} and \mathfrak{B} are given by

$$\begin{aligned} \mathfrak{A} &= \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha \left(\frac{x}{y} \right)^{\beta p} f^p(x) dx dy, \\ \mathfrak{B} &= \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha \left(\frac{y}{x} \right)^{\beta q} g^q(y) dx dy. \end{aligned}$$

Let us now examine the exact expressions for \mathfrak{A} and \mathfrak{B} sequentially.

Applying the Fubini–Tonelli integral theorem (which is possible because the integrand is non-negative), we can express \mathfrak{A} as

$$\mathfrak{A} = \int_{(0,+\infty)} x f^p(x) \left\{ \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha \left(\frac{x}{y} \right)^{\beta p} \frac{1}{x} dy \right\} dx.$$

Let us now concentrate on the integral term with respect to y . Making the change of variables $u = y/x$ and applying standard integration techniques, together with the condition $\alpha > |\beta p - 1|$ (which implies $\alpha - \beta p + 1 > 0$ and $-\alpha - \beta p + 1 < 0$), we obtain

$$\begin{aligned} & \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha \left(\frac{x}{y} \right)^{\beta p} \frac{1}{x} dy = \int_{(0,+\infty)} \left[\min \left(\frac{1}{u}, u \right) \right]^\alpha \frac{1}{u^{\beta p}} du \\ &= \int_{(0,1)} \left[\min \left(\frac{1}{u}, u \right) \right]^\alpha \frac{1}{u^{\beta p}} du + \int_{(1,+\infty)} \left[\min \left(\frac{1}{u}, u \right) \right]^\alpha \frac{1}{u^{\beta p}} du \\ &= \int_{(0,1)} u^\alpha \times \frac{1}{u^{\beta p}} du + \int_{(1,+\infty)} \frac{1}{u^\alpha} \times \frac{1}{u^{\beta p}} du \\ &= \int_{(0,1)} u^{\alpha-\beta p} du + \int_{(1,+\infty)} u^{-\alpha-\beta p} du \\ &= \left[\frac{1}{\alpha - \beta p + 1} u^{\alpha-\beta p+1} \right]_{(0,1)} + \left[\frac{1}{-\alpha - \beta p + 1} u^{-\alpha-\beta p+1} \right]_{(1,+\infty)} \\ &= \frac{1}{\alpha - \beta p + 1} + \frac{1}{\alpha + \beta p - 1} \\ &= \frac{2\alpha}{(\alpha - \beta p + 1)(\alpha + \beta p - 1)}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \mathfrak{A} &= \int_{(0,+\infty)} x f^p(x) \frac{2\alpha}{(\alpha - \beta p + 1)(\alpha + \beta p - 1)} dx \\ &= \frac{2\alpha}{(\alpha - \beta p + 1)(\alpha + \beta p - 1)} \int_{(0,+\infty)} x f^p(x) dx. \end{aligned} \tag{2}$$

For \mathfrak{B} , proceeding analogously, we obtain

$$\begin{aligned} \mathfrak{B} &= \int_{(0,+\infty)} y g^q(y) \frac{2\alpha}{(\alpha - \beta q + 1)(\alpha + \beta q - 1)} dy \\ &= \frac{2\alpha}{(\alpha - \beta q + 1)(\alpha + \beta q - 1)} \int_{(0,+\infty)} y g^q(y) dy. \end{aligned} \tag{3}$$

Joining Equations (1), (2) and (3), we get

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x)g(y)dx dy \\ & \leq \left[\frac{2\alpha}{(\alpha - \beta p + 1)(\alpha + \beta p - 1)} \int_{(0,+\infty)} x f^p(x) dx \right]^{1/p} \\ & \quad \times \left[\frac{2\alpha}{(\alpha - \beta q + 1)(\alpha + \beta q - 1)} \int_{(0,+\infty)} y g^q(y) dy \right]^{1/q} \\ & = \frac{2\alpha}{(\alpha - \beta p + 1)^{1/p}(\alpha + \beta p - 1)^{1/p}(\alpha - \beta q + 1)^{1/q}(\alpha + \beta q - 1)^{1/q}} \\ & \quad \times \left[\int_{(0,+\infty)} x f^p(x) dx \right]^{1/p} \left[\int_{(0,+\infty)} y g^q(y) dy \right]^{1/q}. \end{aligned}$$

This concludes the proof of Theorem 2.1. □

Remark 2.2. Using the following classical minimum formula: $\min(a, b) = (1/2)[a + b - |a - b|]$, with $a, b \in \mathbb{R}$, another formulation of the inequality in Theorem 2.1 is

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\frac{x}{y} + \frac{y}{x} - \left| \frac{x}{y} - \frac{y}{x} \right| \right]^\alpha f(x)g(y)dx dy \\ & \leq \frac{2^{\alpha+1}\alpha}{(\alpha - \beta p + 1)^{1/p}(\alpha + \beta p - 1)^{1/p}(\alpha - \beta q + 1)^{1/q}(\alpha + \beta q - 1)^{1/q}} \\ & \quad \times \left[\int_{(0,+\infty)} x f^p(x) dx \right]^{1/p} \left[\int_{(0,+\infty)} y g^q(y) dy \right]^{1/q}. \end{aligned}$$

Remark 2.3. If we take $p = 2$ (so $q = 2$), under the condition $\alpha > |2\beta - 1|$, the inequality in Theorem 2.1 becomes

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x)g(y)dx dy \\ & \leq \frac{2\alpha}{\alpha^2 - (2\beta - 1)^2} \left[\int_{(0,+\infty)} x f^2(x) dx \right]^{1/2} \left[\int_{(0,+\infty)} y g^2(y) dy \right]^{1/2}. \end{aligned}$$

To the best of our knowledge, Theorem 2.1 introduces a new minimum fractional Hardy–Hilbert-type integral inequality to the literature. We emphasize its flexibility, which comes from the adjustable parameters α and β , as well as the clear and tractable condition $\alpha > \max(|\beta p - 1|, |\beta q - 1|)$. Several related results derived from this theorem constitute the focus of the remainder of the article.

3 Secondary results

The theorem below establishes an integral inequality involving a single function, obtained as a direct consequence of Theorem 2.1.

Theorem 3.1. Let $p > 1, q = p/(p - 1), \alpha, \beta > 0$ such that $\alpha > \max(|\beta p - 1|, |\beta q - 1|)$, and $f, g : (0, +\infty) \mapsto (0, +\infty)$ be two functions. Then we have

$$\int_{(0,+\infty)} y^{-(p-1)} \left[\int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right]^p dy \leq \frac{(2\alpha)^p}{(\alpha - \beta p + 1)(\alpha + \beta p - 1)(\alpha - \beta q + 1)^{p-1}(\alpha + \beta q - 1)^{p-1}} \int_{(0,+\infty)} x f^p(x) dx$$

provided that the integral involved in the upper bound converges.

Proof. Let us set

$$\mathfrak{C} = \int_{(0,+\infty)} y^{-(p-1)} \left[\int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right]^p dy.$$

We can write

$$\begin{aligned} \mathfrak{C} &= \int_{(0,+\infty)} \left[\int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right] \left[y^{-1} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right]^{p-1} dy \\ &= \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) g_{\dagger}(y) dx dy \end{aligned} \tag{4}$$

where

$$g_{\dagger}(y) = \left[y^{-1} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right]^{p-1}.$$

Applying Theorem 2.1 to f and g_{\dagger} , we obtain

$$\begin{aligned} &\int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) g_{\dagger}(y) dx dy \\ &\leq \frac{2\alpha}{(\alpha - \beta p + 1)^{1/p}(\alpha + \beta p - 1)^{1/p}(\alpha - \beta q + 1)^{1/q}(\alpha + \beta q - 1)^{1/q}} \\ &\quad \times \left[\int_{(0,+\infty)} x f^p(x) dx \right]^{1/p} \left[\int_{(0,+\infty)} y g_{\dagger}^q(y) dy \right]^{1/q}. \end{aligned} \tag{5}$$

Moreover, using $q(p - 1) = p$, we have

$$\begin{aligned} &\int_{(0,+\infty)} y g_{\dagger}^q(y) dy = \int_{(0,+\infty)} y \left[y^{-1} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right]^{q(p-1)} dy \\ &= \int_{(0,+\infty)} y \left[y^{-1} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right]^p dy \\ &= \int_{(0,+\infty)} y^{-(p-1)} \left[\int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right]^p dy \\ &= \mathfrak{C}. \end{aligned} \tag{6}$$

Joining Equations (4), (5) and (6), we get

$$\mathfrak{C} \leq \frac{2\alpha}{(\alpha - \beta p + 1)^{1/p}(\alpha + \beta p - 1)^{1/p}(\alpha - \beta q + 1)^{1/q}(\alpha + \beta q - 1)^{1/q}} \times \left[\int_{(0,+\infty)} x f^p(x) dx \right]^{1/p} \mathfrak{C}^{1/q}.$$

Using $1 - 1/q = 1/p$, we obtain

$$\mathfrak{C}^{1/p} \leq \frac{2\alpha}{(\alpha - \beta p + 1)^{1/p}(\alpha + \beta p - 1)^{1/p}(\alpha - \beta q + 1)^{1/q}(\alpha + \beta q - 1)^{1/q}} \times \left[\int_{(0,+\infty)} x f^p(x) dx \right]^{1/p}.$$

Raising both sides at the exponent p , and taking into account the definition of \mathfrak{C} and that $p/q = p - 1$, we obtain

$$\int_{(0,+\infty)} y^{-(p-1)} \left[\int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right]^p dy \leq \frac{(2\alpha)^p}{(\alpha - \beta p + 1)(\alpha + \beta p - 1)(\alpha - \beta q + 1)^{p-1}(\alpha + \beta q - 1)^{p-1}} \int_{(0,+\infty)} x f^p(x) dx.$$

This completes the proof of Theorem 3.1. □

Remark 3.2. If we take $p = 2$, under the condition $\alpha > |2\beta - 1|$, the inequality in Theorem 3.1 becomes

$$\int_{(0,+\infty)} y^{-1} \left[\int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f(x) dx \right]^2 dy \leq \frac{(2\alpha)^2}{[\alpha^2 - (2\beta - 1)^2]^2} \int_{(0,+\infty)} x f^2(x) dx.$$

The theorem below is analogous to Theorem 2.1, but it involves the primitives of the main functions. The resulting bound is expressed in terms of the unweighted integral norms of these functions. It is obtained as a consequence of Theorem 2.1 combined with the classical Hardy integral inequality (see [6]).

Theorem 3.3. Let $p > 1$, $q = p/(p - 1)$, $\alpha, \beta > 0$ such that $\alpha > \max(|\beta p - 1|, |\beta q - 1|)$, $f, g : (0, +\infty) \mapsto (0, +\infty)$ be two functions, and, for any $x > 0$,

$$F(x) = \int_0^x f(t) dt, \quad G(x) = \int_0^x g(t) dt,$$

provided that the two integrals converge. Then we have

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha x^{-1-1/p} F(x) y^{-1-1/q} G(y) dx dy \\ & \leq \frac{2\alpha p^2}{(\alpha - \beta p + 1)^{1/p} (\alpha + \beta p - 1)^{1/p} (\alpha - \beta q + 1)^{1/q} (\alpha + \beta q - 1)^{1/q} (p - 1)} \\ & \quad \times \left[\int_{(0,+\infty)} f^p(x) dx \right]^{1/p} \left[\int_{(0,+\infty)} g^q(y) dy \right]^{1/q} \end{aligned}$$

provided that the two integrals involved in the upper bound converge.

Proof. We can write

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha x^{-1-1/p} F(x) y^{-1-1/q} G(y) dx dy \\ & = \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f_\diamond(x) g_\diamond(y) dx dy \end{aligned} \tag{7}$$

where

$$f_\diamond(x) = x^{-1-1/p} F(x), \quad g_\diamond(y) = y^{-1-1/q} G(y).$$

Applying Theorem 2.1 to f_\diamond and g_\diamond , we obtain

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha f_\diamond(x) g_\diamond(y) dx dy \\ & \leq \frac{2\alpha}{(\alpha - \beta p + 1)^{1/p} (\alpha + \beta p - 1)^{1/p} (\alpha - \beta q + 1)^{1/q} (\alpha + \beta q - 1)^{1/q}} \\ & \quad \times \left[\int_{(0,+\infty)} x f_\diamond^p(x) dx \right]^{1/p} \left[\int_{(0,+\infty)} y g_\diamond^q(y) dy \right]^{1/q}. \end{aligned} \tag{8}$$

Applying the classical Hardy integral inequality to f , we obtain

$$\begin{aligned} & \int_{(0,+\infty)} x f_\diamond^p(x) dx = \int_{(0,+\infty)} x [x^{-1-1/p} F(x)]^p dx = \int_{(0,+\infty)} x^{-p} F^p(x) dx \\ & \leq \left(\frac{p}{p-1} \right)^p \int_{(0,+\infty)} f^p(x) dx. \end{aligned} \tag{9}$$

Similarly, using $p = q/(q - 1)$, we get

$$\begin{aligned} & \int_{(0,+\infty)} y g_\diamond^q(y) dy = \int_{(0,+\infty)} y [y^{-1-1/q} G(y)]^q dy = \int_{(0,+\infty)} y^{-q} G^q(y) dy \\ & \leq \left(\frac{q}{q-1} \right)^q \int_{(0,+\infty)} g^q(y) dy = p^q \int_{(0,+\infty)} g^q(y) dy. \end{aligned} \tag{10}$$

Joining Equations (7), (8), (9) and (10), we find that

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha x^{-1-1/p} F(x) y^{-1-1/q} G(y) dx dy \\ & \leq \frac{2\alpha}{(\alpha - \beta p + 1)^{1/p} (\alpha + \beta p - 1)^{1/p} (\alpha - \beta q + 1)^{1/q} (\alpha + \beta q - 1)^{1/q}} \\ & \quad \times \left[\left(\frac{p}{p-1} \right)^p \int_{(0,+\infty)} f^p(x) dx \right]^{1/p} \left[p^q \int_{(0,+\infty)} g^q(y) dy \right]^{1/q} \\ & = \frac{2\alpha p^2}{(\alpha - \beta p + 1)^{1/p} (\alpha + \beta p - 1)^{1/p} (\alpha - \beta q + 1)^{1/q} (\alpha + \beta q - 1)^{1/q} (p-1)} \\ & \quad \times \left[\int_{(0,+\infty)} f^p(x) dx \right]^{1/p} \left[\int_{(0,+\infty)} g^q(y) dy \right]^{1/q}. \end{aligned}$$

This concludes the proof of Theorem 3.3. □

Remark 3.4. If we take $p = 2$, under the condition $\alpha > |2\beta - 1|$, the inequality in Theorem 3.3 becomes

$$\begin{aligned} & \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x} \right) \right]^\alpha x^{-3/2} F(x) y^{-3/2} G(y) dx dy \\ & \leq \frac{8\alpha}{\alpha^2 - (2\beta - 1)^2} \left[\int_{(0,+\infty)} f^2(x) dx \right]^{1/2} \left[\int_{(0,+\infty)} g^2(y) dy \right]^{1/2}. \end{aligned}$$

To the best of our knowledge, like Theorem 2.1, Theorem 3.3 introduces another new Hardy–Hilbert-type integral inequality to the literature.

4 Conclusion

In this article, we have established a new minimum fractional Hardy–Hilbert-type integral inequality. It provides a sharp bound that depends on a minimum function of the ratios x/y and y/x as well as a key parameter, α , and two main functions, f and g . Building on this main theorem, several secondary inequalities were derived, highlighting the versatility of the approach and its potential for further generalizations. Future work could explore multidimensional extensions, such as a sharp upper bound for a triple integral of the form

$$\int_{(0,+\infty)} \int_{(0,+\infty)} \int_{(0,+\infty)} \left[\min \left(\frac{x}{y}, \frac{y}{x}, \frac{x}{z}, \frac{z}{x}, \frac{y}{z}, \frac{z}{y} \right) \right]^\alpha f(x)g(y)h(z) dx dy dz.$$

In addition, we could consider applications in operator theory and its connections with fractional differential equations and other areas of functional analysis.

References

- [1] Laith Emil Azar. On some extensions of Hardy–Hilbert’s inequality and applications. *Journal of Inequalities and Applications*, 2008:546829, 2008. DOI: 10.1155/2008/546829.
- [2] Laith Emil Azar, Tserendorj Batbold, Nizar Kh. Al-Oushoush, and Mario Krnić. A class of more accurate Hilbert-type inequalities involving partial sums. *Journal of Inequalities and Special Functions*, 15(1):1–10, 2024. DOI: 10.54379/jiasf-2024-1-1
- [3] Tserendorj Batbold, Mario Krnić, and Predrag Vuković. A unified approach to fractal Hilbert-type inequalities. *Journal of Inequalities and Applications*, 2019:117, 2019. DOI: 10.1186/s13660-019-2076-9
- [4] Qiang Chen and Bicheng Yang. A survey on the study of Hilbert-type inequalities. *Journal of Inequalities and Applications*, 2015:302, 2015. DOI: 10.1186/s13660-015-0829-7.
- [5] Christophe Chesneau. Refining and extending two special Hardy–Hilbert-type integral inequalities. *Annals of Mathematics and Computer Science*, 28:21–45, 2025. DOI: 10.56947/amcs.v28.513.
- [6] Godfrey Harold Hardy, John Edensor Littlewood, George Pólya. *Inequalities*. Cambridge University Press, 1934. ISBN: 9780521358804.
- [7] Mario Krnić and Predrag Vuković. Multidimensional Hilbert-type inequalities obtained via local fractional calculus. *Acta Applicandae Mathematicae*, 169:667–680, 2020. DOI: 10.1007/s10440-020-00317-x.
- [8] Yongjin Li and Bing He. On inequalities of Hilbert’s type. *Bulletin of the Australian Mathematical Society*, 76(1):1–13, 2007. DOI: 10.1017/S0004972700039423.
- [9] Yongjin Li, Jing Wu, and Bing He. A new Hilbert-type integral inequality and the equivalent form. *International Journal of Mathematics and Mathematical Sciences*, 2006(45378):1–6, 2006. DOI: 10.1155/IJMMS/2006/45378.
- [10] Michael Th. Rassias and Bicheng Yang. A multidimensional Hilbert-type integral inequality related to the Riemann zeta function. In *Applications of Mathematics and Informatics in Science and Engineering*, volume 91 of *Optimization and Its Applications*, pages 417–433. Springer, 2014. DOI: 10.1007/978-3-319-04720-1_26.
- [11] Michael Th. Rassias and Bicheng Yang. Equivalent properties of a Hilbert-type integral inequality with the best constant factor related to the Hurwitz zeta function. *Annals of Functional Analysis*, 9(2):282–295, 2018. DOI: 10.1215/20088752-2017-0031.
- [12] Michael Th. Rassias and Bicheng Yang. On a Hilbert-type integral inequality in the whole plane related to the extended Riemann zeta function. *Complex Analysis and Operator Theory*, 13:1765–1782, 2019. DOI: 10.1007/s11785-018-0830-5.

- [13] Aziz Saglam, Hüseyin Yildirim, and Mehmet Zeki Sarikaya. Generalization of Hardy–Hilbert’s inequality and applications. *Kyungpook Mathematical Journal*, 50(1):131–152, 2010. DOI: 10.5666/KMJ.2010.50.1.131.
- [14] Mehmet Zeki Sarikaya and Mehmet Sabir Bingöl. Recent developments of integral inequalities of the Hardy–Hilbert type. *Turkish Journal of Inequalities*, 8(2):43–54, 2024.
- [15] Waadallah Tawfeeq Sulaiman. New Hardy–Hilbert’s-type integral inequalities. *International Mathematical Forum*, 3(43):2139–2147, 2008.
- [16] Waadallah Tawfeeq Sulaiman. An extension of Hardy–Hilbert’s integral inequality. *African Diaspora Journal of Mathematics*, 10(2):66–71, 2010.
- [17] Waadallah Tawfeeq Sulaiman. Hardy-Hilbert’s integral inequality in new kinds. *Mathematical Communications*, 15(2):453–461, 2010.
- [18] Waadallah Tawfeeq Sulaiman. New kinds of Hardy–Hilbert’s integral inequalities. *Applied Mathematics Letters*, 23(4):361–365, 2010. DOI: 10.1016/j.aml.2009.10.011.
- [19] Bajou Sun. Best generalization of a Hilbert type inequality. *Journal of Inequalities in Pure and Applied Mathematics*, 7(3):113, 2006.
- [20] Bicheng Yang. *Hilbert-Type Integral Inequalities*. Bentham Books, 2009. ISBN: 9781608050338.
- [21] Bicheng Yang and Michael Th. Rassias. *On Hilbert-type and Hardy-type integral inequalities and applications*. Volume of *Briefs in Mathematics*. Springer, 2019. DOI: 10.1007/978-3-030-29268-3.
- [22] Bicheng Yang and Michael Th. Rassias. *On extended Hardy–Hilbert integral inequalities and applications*. World Scientific, 2023. DOI: 10.1142/13164.